arXiv:hep-ph/9806297v1 6 Jun 1998

THE TENSOR STRUCTURE FUNCTION $b_2(x, Q^2)$ OF THE DEUTERON AT SMALL x

W. SCHÄFER
Institut für Kernphysik, Forschungszentrum Jülich, D-52425 Jülich

The dependence of the diffractive nuclear shadowing and the nuclear pion excess effect on the deuteron spin alignment leads to a substantial tensor polarisation of sea partons in the deuteron. The corresponding tensor structure function $b_2(x, Q^2)$ rises towards small x, and we predict the about one percent tensor asymmetry $A_2(x, Q^2) = b_2(x, Q^2)/F_{2D}(x, Q^2)$. We calculate the shadowing of the gluon distribution in the deuteron for two different pomeron gluon-distributions. For a hard gluon dominated $\mathbb{P}$-structure function, as advocated by H1, we find large gluon shadowing, which appears to be in conflict with the DGLAP analysis of nuclear SF's.

1 The shadowing driven tensor structure function

It is a common theoretical prejudice, that in the high energy limit ($x \to 0$ in DIS), total cross sections cease to depend upon the beam and target polarizations. However we shall see in the following, that the tensor polarization of sea quarks in a deuteron is substantial and even rises for $x \to 0$ [1]. Let $\vec{S}$ be a spin-1 operator, then the *tensor polarization* $T_{ik} = \frac{1}{2}(S_i S_k + S_k S_i - \frac{2}{3}\vec{S}^2 \delta_{ik})$ is a T-even and P-even observable. As such it is accessable in DIS of *unpolarized* leptons on the *(tensor-)polarized* deuteron. Let $\vec{n}$ be the deuteron-spin quantization axis, and $\lambda = \vec{S} \cdot \vec{n}$ the spin projection upon the quantization axis. Then the total photoabsorption cross sections $\sigma^{(\lambda)}(\gamma^*_{T,L} D)$ bear a dependence upon the deuteron polarization state, and thus on the tensor polarization [2] (for the case of real photons see also[3]): $\sigma^{(\lambda)}(\gamma^*_{T,L} D) = \sigma_0 + \sigma_2 T_{ik} n_i n_k + ...$ The corresponding structure functions are introduced through

$$F_2^{(\lambda)} = \frac{Q^2}{4\pi^2 \alpha_{em}} (\sigma^{(\lambda)}(\gamma^*_T D) + \sigma^{(\lambda)}(\gamma^*_L D)). \quad (1)$$

If one choses the $\gamma^* - D$-collision axis for the quantization axis ($z-$ axis), the structure function b_2 can be defined as:

$$\begin{aligned} b_2(x, Q^2) &= \frac{1}{2}[F_2^{(+)}(x, Q^2) + F_2^{(-)}(x, Q^2) - 2F_2^{(0)}(x, Q^2)] \\ &= \frac{1}{2} \sum_f e_f^2 x [q_f^{(+)}(x, Q^2) + q_f^{(-)}(x, Q^2) - 2q_f^{(0)}(x, Q^2)]. \end{aligned} \quad (2)$$

Here the last line is the expression for b_2 in the parton model, and $q_f^{(\lambda)}$ stands for the parton density in the hadron with spin projection λ on the $z-$axis. In the same manner one defines the transverse $b_T = 2xb_1$ and longitudinal $b_L = b_2 - b_T$ structure functions. It was only the impulse approximation (IA) that had been employed in the discussion of the tensor structure functions before [2,4]. What has been found is a contribution to b_2 that vanishes in the limit $x \to 0$ and is only the per-mille effect even at moderate x. Furthermore, in the IA one can derive the sum rule $\int_0^1 dx\ b_1(x, Q^2) = 0$. A conjecture had been made by Close and Kumano [5], for it to hold also beyond the impulse approximation, under the assumption of an unpolarised sea. We shall see that the results of the beyond-the-IA evaluation of the tensor structure function imply that such a sum rule for b_1 cannot exist. The appropriate framework for DIS on nuclear targets is the Glauber-Gribov multiple scattering theory[6,8]. The inelastic shadowing diagrams lead to the nuclear eclipse effect: $\sigma_{tot}(\gamma^* D) = \sigma_{tot}(\gamma^* p) + \sigma_{tot}(\gamma^* n) - \Delta\sigma_{sh}(\gamma^* D)$ and two production mechanisms of the inelastically exited intermediate state X have to be distinguished: in diffractive nuclear shadowing (NSH) the state X is produced by pomeron exchange, the second mechanism is the exitation of X by pion exchange. The diffractive NSH contribution can readily be calculated using Gribov's theory extended to DIS in[8]:



Figure 1: Impulse approximation (a) and Gribov's inelastic shadowing (b),(c) diagrams for DIS on the deuteron.

$$\Delta F_{sh}^{(\lambda)}(x, Q^2) = \frac{2}{\pi} \frac{Q^2}{4\pi^2 \alpha_e m} \int d^2\vec{k}_\perp \int dM^2 S_D^{(\lambda)}(4\vec{k}^2) \frac{d\sigma^D(\gamma^* \to X)}{dt dM^2}\bigg|_{t=-\vec{k}^2} . \quad (3)$$

Here M is the mass of the state X, $\vec{k} = (\vec{k}_\perp, k_z)$ is the momentum of the pomeron, and $S_D^{(\lambda)}(4\vec{k}^2) = \int d^3\vec{r} |\Psi_D^{(\lambda)}(\vec{r})|^2 exp(-i\vec{r}\cdot\vec{k})$ denotes the deuteron formfactor. It is of crucial importance to include in the calculation the $D-$wave admixture in the deuteron wavefunction $\Psi_D^{(\lambda)}$, as it turns out that the diffractive NSH contribution to the tensor structure function is mainly due to the $S-D-$ interference. A further ingredient of the calculation is the differential cross section $d\sigma^D(\gamma^* \to X)/dtdM^2$ for the diffraction dissociation of the virtual photon into the state X on the proton, which can be written in terms of the diffractive structure function. We make use of a theoretical calculation of the diffractive cross section [7] that reproduces the HERA data on diffraction – this should be demanded of any reliable calculation of the nuclear shadowing. Schematically, the diffractive contribution can be cast into the convolution form:

$$\Delta F_{sh}^{(\lambda)}(x,Q^2) = \int_x^1 \frac{dx_{\mathbf{IP}}}{x_{\mathbf{IP}}} \Delta n_{\mathbf{IP}/D}^{(\lambda)}(x_{\mathbf{IP}}) F_{2\mathbf{IP}}(\frac{x}{x_{\mathbf{IP}}},Q^2), \tag{4}$$

where $\Delta n_{\mathbf{IP}/D}^{(\lambda)}(x_{\mathbf{IP}})$ is the nuclear modification of the pomeron flux. The second mechanism is pion exchange: it dominates $\gamma^* p \to nX$, and contributes substantially to $\gamma^* p \to pX$ at moderately small $x_{\mathbf{IP}}$. One can derive an analogous convolution formula as for the pomeron contribution, only the pomeron flux and structure function get replaced by the corresponding pionic quantities. From these representations one can read off the evolution properties of the two contributions: diffractive NSH is known to be the leading twist, DGLAP-evolving effect [8]. That the pionic contribution is DGLAP evolving is self-evident, as the Q^2-dependence enters only through the pion structure function. Our results for b_2 and the related asymmetry $A_2 = b_2/F_{2D}$ can be seen in fig.2a,b) .

2 Constraining the glue in the Pomeron by nuclear shadowing

At present, there are several conflicting claims about the gluon density in the pomeron: the colour dipole model calculations by Nikolaev and Zakharov, on which our evaluations of the diffractive NSH are based predicts that gluons and quarks carry about an equal fraction of the pomeron's momentum, $\langle x_G \rangle \simeq \langle x_v \rangle$. In contrast to that the DGLAP fits to the H1 diffractive structure function data prefer the gluon dominated pomeron: $\langle x_G \rangle \gg \langle x_v \rangle$. The two options for glue in the pomeron, $G_{\mathbf{IP}}(x,Q^2) = xg_{\mathbf{IP}}(x,Q^2)$ give drastically different predictions for the shadowing correction

$$\Delta G_{sh}^{(\lambda)}(x,Q^2) = \int_x^1 \frac{dx_{\mathbf{IP}}}{x_{\mathbf{IP}}} \Delta n_{\mathbf{IP}/D}^{(\lambda)}(x_{\mathbf{IP}}) G_{\mathbf{IP}}(\frac{x}{x_{\mathbf{IP}}},Q^2), \tag{5}$$

to the gluon structure function of the deuteron, see fig 2c). A similar formula holds for the pionic contribution. There are no direct experimental data on nuclear shadowing of glue in the deuteron, but precision data on shadowing in DIS on heavy nuclei allow one to perform the DGLAP analysis of nuclear structure functions. Eskola et al. in their recent analysis find



Figure 2: (a) The tensor structure function of the deuteron $b_2(x, Q^2)$ at $Q^2 = 10 GeV^2$ and its diffractive NSH and pion excess components. (b) the tensor polarization $A_2(x, Q^2)$ of quarks and antiquarks in the deuteron at $Q^2 = 10 GeV^2$. Also shown (by the dotted line) is the longitudinal tensor structure function $b_L(x, Q^2)$ (in (a)) and the corresponding asymmetry (in(b)).(c) Diffractive shadowing correction to the gluon distribution in the deuteron for two different pomeron gluon distributions: the H1 parametrization with $\langle x_G \rangle \gg \langle x_v \rangle$ and the prediction from NZ, for which $\langle x_G \rangle \simeq \langle x_v \rangle$. The dashed lines are for $Q^2 = 4 GeV^2$, the solid ones for $Q^2 = 2.25 GeV^2$

$R_G^A = \Delta G_{sh}^{(A)}(x, Q_0^2)/AG(x, Q_0^2) \simeq 0.15$ at $Q_0^2 \simeq 2.25 GeV^2, x \simeq 0.001$ for the ^{12}C–nucleus which is about the same as shadowing of sea quarks. The straightforward nuclear size arguments suggest[10] that $R_G^A(deuteron) \simeq 0.3 \cdot R_G^A(^4He) \simeq 0.2 \cdot R_G^A(^{12}C)$, i.e. a $\sim 3\%$ effect for the deuteron, which agrees with the result for the NZ-glue in the pomeron. In contrast, the very strong shadowing predicted by the H1 glue is in an obvious conflict with the NMC experimental data on nuclear shadowing.

3 Conclusions

We have demonstrated that the deuteron tensor structure function at small x is driven by the inelastic nuclear shadowing. It is rising towards small x, which makes it a unique spin effect – all other spin effects in total cross sections are known to die out at high energies. It was shown that this fact leads to the conclusion, that the Close-Kumano sum-rule does not exist. Our findings of a rise of b_2 at small x have also been confirmed by calculations of other groups[13]. Further it was demonstrated, that a hard gluon dominated pomeron structure function leads to strong gluon shadowing in the deuteron, which is quite suspect in view of the DGLAP-phenomenology of nuclear structure functions.



Figure 3: (a)Shadowing contributions to the deuteron gluon SF: diffractive NSH(solid line) and pionic contribution (dashed line).(b)Gluon tensor asymmetry $A_G(x,Q^2) = (G^{(\pm)}(x,Q^2) - G^{(0)}(x,Q^2))/G^D(x,Q^2)$: diffractive (solid line) and pionic (dashed line) contributions; $Q^2 = 4GeV^2$.

References

1. N.N. Nikolaev and W.Schäfer, *Phys. Lett.* B **398**, 245 (1997)
2. P. Hoodbhoy,R.L. Jaffe and A. Manohar, *Nucl. Phys.* B **312**, 571 (1988)
3. A. Pais, *Phys. Rev. Lett.* **19**, 544 (1967)
4. A.Yu. Umnikov *Phys. Lett.* B **391**, 177 (1997)
5. F. Close and S. Kumano, *Phys. Rev.* D **42**, 2377 (1990).
6. V.N. Gribov, *Sov. Phys. JETP* **29**, 483 (1969).
7. M. Genovese, N.N. Nikolaev and B.G. Zakharov, *Phys. Lett.* B **378**, 347 (1996).
8. N.N. Nikolaev and B.G. Zakharov, *Z. Phys.* C **49**, 607 (1991)
9. N.N. Nikolaev and B.G. Zakharov, *Z. Phys.* C **53**, 331 (1992)
10. V. Barone, M. Genovese, N.N. Nikolaev, E. Predazzi and B.G. Zakharov, *Z. Phys.* C **58**, 541 (1993)
11. K.J. Eskola et al., e-print-archives hep-ph/9802350
12. H1 collab.:C. Adloff et al., *Z. Phys.* C **76**, 613 (1997)
13. J. Edelmann, G. Piller and W. Weise, e-print-archives hep-ph/9709455, K. Bora and R. Jaffe, e-print-archives hep-ph/9711323